Skill Procedures: Nebulized Albuterol Administration

I. Usage

1. This protocol describes how to administer albuterol by means of an oxygen nebulizer.

II. Procedural Protocols

- 1. Assemble equipment.
- 2. Check medication (2.5mg albuterol in 3cc normal saline) and empty into nebulizer chamber.
- 3. Attach tubing to an oxygen outlet flowing at 10 liters per minute.



- 4. Administer solution by having patient put lips around mouth piece and breath deeply. For children unable to comply, hold mouth piece near nose and mouth.
 - A. For patient's who have difficulty using the mouthpiece, it is permissible to attach the nebulizer to a simple face mask.
 - i. Monitor patient closely during the nebulizer treatment.
 - ii. Increase oxygen flow upon completion of treatment if face mask is still utilized.



- 5. Monitor patients vital signs and lung sounds. Contact On Line Medical Control if:
 - A. Pulse rate increases by 30 or more beats per minute.
 - B. Patient develops PVC's (or ventricular ectopy increases).
 - C. Respiratory distress increases.
- 6. Monitor and report patient's response to medication.
- 7. Record medication given, dose, amount and time on the Patient Care Report.

Effective Date: 1 May 2002 Revision Number: N/A
Revision Date: N/A Page J15.1

Skill Procedures: Nebulized Albuterol Administration

This Page Intentionally Left Blank

Effective Date: 1 May 2002 Revision Number: N/A
Revision Date: N/A Page J15.2